

MR. ZHENG SHI (Orcid ID : 0000-0002-5067-9977)

DR. KEVIN RORY WILCOX (Orcid ID : 0000-0001-6829-1148)

DR. LARA SOUZA (Orcid ID : 0000-0001-6005-8667)

DR. JIANG JIANG (Orcid ID : 0000-0001-5058-8664)

MR. CHANG GYO JUNG (Orcid ID : 0000-0002-9845-7732)

Article type : Primary Research Articles

Successional change in species composition alters climate sensitivity of grassland productivity

Running title: Response of grassland productivity to climate change

Zheng Shi^{1,2*}, Yang Lin³, Kevin R. Wilcox², Lara Souza², Lifen Jiang⁴, Jiang Jiang⁵, Chang Gyo Jung⁴, Xia Xu¹, Mengting Yuan^{2,6}, Xue Guo^{2,7}, Liyou Wu^{2,6}, Jizhong Zhou^{2,6,8,9}, and Yiqi Luo⁴

¹Co-Innovation Center for Sustainable Forestry in Southern China, College of Biology and the Environment, Nanjing Forestry University, Nanjing, 210037, China

²Department of Microbiology & Plant Biology, University of Oklahoma, Norman, OK 73019, USA

³Department of Environmental Science, Policy, and Management, University of California, Berkeley, CA 94720, USA

⁴Center for Ecosystem Science and Society, Northern Arizona University, Flagstaff, AZ 86011, USA

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/gcb.14333

This article is protected by copyright. All rights reserved.

⁵Key Laboratory of Soil and Water Conservation and Ecological Restoration in Jiangsu Province, Collaborative Innovation Center of Sustainable Forestry in Southern China of Jiangsu Province, Nanjing Forestry University, Nanjing, 210037, China

⁶Institute for Environmental Genomics, University of Oklahoma, Norman, OK 73019, USA.

⁷ School of Minerals Processing and Bioengineering, Central South University, Changsha 410083, China

⁸State Key Joint Laboratory of Environment Simulation and Pollution Control, School of Environment, Tsinghua University, Beijing, 100084, PR China.

⁹Earth and Environmental Sciences, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA.

*: Authors for correspondence: Zheng Shi (zheng.shi@ou.edu)

Mailing address: 101 David L Boren Blvd., Norman, Oklahoma 73072, USA

Tel.: +1 405-325-7619 Fax: +1 405-325-1595

Keywords: succession; anthropogenic perturbation; biotic competition; long-term experiment; manipulative experiment

Author contributions: Y.L. designed the experiments. Z.S., K.W., L.J., J.J., C.J., X.X., M.Y., and X.G. collected the data. Z.S., Y.L., and K.W. performed data analyses. All authors contributed to the writing and discussions. **Competing interests:** The authors declare that they have no competing interests.

Abstract

Succession theory predicts altered sensitivity of ecosystem functions to disturbance (i.e., climate change) due to the temporal shift in plant community composition. However, empirical evidence in global change experiments is lacking to support this prediction. Here, we present findings from an eight-year long-term global change experiment with warming and altered precipitation manipulation (double and halved amount). First, we observed a temporal shift in species composition over eight years, resulting in a transition from an annual C₃-dominant plant community to a perennial C₄-dominant plant community. This successional transition was independent of any experimental treatments. During the successional transition, the response of aboveground net primary productivity (ANPP) to precipitation addition magnified from neutral to +45.3%, while the response to halved precipitation attenuated substantially from -17.6% to neutral. However, warming did not affect ANPP in either state. The findings further reveal that the time-dependent climate sensitivity may be regulated by successional change in species composition, highlighting the importance of vegetation dynamics in regulating the response of ecosystem productivity to precipitation change.

Introduction

Long-term shift in the sensitivity of ecosystem functions to climatic change (Smith *et al.*, 2009) has been widely supported with empirical evidence (Fu *et al.*, 2015, Melillo *et al.*, 2002, Piao *et al.*, 2017, Wu *et al.*, 2012, Yvon-Durocher *et al.*, 2017). Multiple mechanisms have been proposed for causing the altered climate sensitivity of ecosystem functions. Among them are organism acclimation (Bradford *et al.*, 2008, Crowther & Bradford, 2013, Smith *et al.*, 2016), substrate depletion (Luo *et al.*, 2001, Melillo *et al.*, 2002), nutrient and species diversity loss (Wu *et al.*, 2012) underlying the dampened sensitivity over time; and

species compositional change responsible for the amplified sensitivity over time (Wilcox *et al.*, 2016, Xu *et al.*, 2015). Ecological theory also predicts that ecosystem responses to perturbation (such as climate change) vary with successional states (Odum, 1969) and dynamic vegetation composition driven by succession may modulate ecosystem responses to climate change.

Cross-site experiments have suggested that plant communities at different successional states respond differently to climate change. Kröel-Dulay *et al.* (2015) showed that plant species richness and composition were responsive to experimental warming or drought in three early-successional European shrublands, but were resistant in mature shrubland ecosystems, along a large climatic gradient. Similarly, Grime *et al.* (2000) reported that plant community biomass and composition in an early-successional fertile limestone grassland showed greater responses to experimental warming and drought than those in a late-successional infertile grassland. These findings suggest that differences in plant functional groups, resulting from differences in species composition, along successional states be a key underlying mechanism for the differential responses to climate change. However, climatic and edaphic variables often co-vary with the successional states in these cross-site experiments, which makes these conclusions and the role of functional groups in regulating climate sensitivity elusive.

Old field succession is a form of secondary, ecological succession occurring on abandoned agricultural land. As the old field succession proceeds, the plant communities differ in several ways. Early successional communities consist of annual weedy grasses and forbs such as ragweed (*Ambrosia*) and crabgrass (*Digitaria*), and the following communities at a later stage are dominated primarily by perennial bunch grasses replacing the annual pioneer species (Booth, 1941, Davidson, 1993). Major mechanisms driving the succession are known to be resource limitation through which changes in nitrogen availability alter species

competition, and aboveground vertebrate grazing (Bardgett & Wardle, 2003, Davidson, 1993, Pickett *et al.*, 1987). In addition, shift in plant functional group (e.g., C₃ and C₄ photosynthetic pathways) may be associated with the successional change due to the possible change in nitrogen availability (Ellery *et al.*, 1992). In contrast, disturbance such as herbivory, grazing and fire may sometimes delay vegetation succession and thus maintain plant community at the earlier successional stage (Davidson, 1993).

Physiological properties of plant functional groups may determine ecosystem responses to climate change. For example, species with C₄ photosynthetic pathways are typically more sensitive to warming and better adapted to dry conditions than species with C₃ photosynthetic pathways (Ehleringer *et al.*, 1997, Lambers *et al.*, 1998). The ecological advantage of C₄ over C₃ species under warmer temperature and drought is due to the fact that C₄ photosynthesis has a highly efficient CO₂ pump that elevates CO₂ concentration in the bundle sheath cell (Lambers *et al.*, 1998). Findings from global change experiments (Hoover *et al.*, 2014, Morgan *et al.*, 2011, Niu *et al.*, 2010, Taylor *et al.*, 2014, Wilcox *et al.*, 2015), isotope data analyses (Ehleringer *et al.*, 1997, Von Fischer *et al.*, 2008) and large-scale model predictions (Cramer *et al.*, 2001, Epstein *et al.*, 2002) have demonstrated that in general (I) climate warming favors C₄ plant species over C₃, (II) drought adversely affects C₃ plant species over C₄ species, and (III) increased precipitation enhances C₄ plant growth to a greater magnitude than C₃ plant growth due to higher photosynthetic water use efficiency (Knapp & Medina, 1999, Lambers *et al.*, 1998). However, the physiological differences between the two functional groups may not translate simply into ecological responses due to the indirect effects by climate change. For example, altered soil moisture by climate change can have an indirect effect on plant growth and therefore may counteract the physiological response of plant functional groups (Mueller *et al.*, 2016); altered plant phenology and biotic competition by climate change may also add another level of complexity to the overall

ecosystem responses (Niu & Wan, 2008, Reyes-Fox *et al.*, 2014, Shi *et al.*, 2015). For example, warming-induced advancement of spring phenology for certain species may have a positive influence on their competitive ability (Parmesan, 2007).

Here, we leverage an eight-year (2009 ~ 2016) global change experiment in a grassland in central Oklahoma, USA where plant dominance of the ecosystem shifted from C₃ to C₄ species after the removal of grazing. This experiment was a factorial manipulation of temperature (+ 3 °C) using infrared heaters and precipitation (halved and double amount) using rainout facility (Xu *et al.*, 2016). Using this unique opportunity, we investigate whether the magnitude of responses of ecosystem productivity to climate change treatments (warming and altered precipitation regime) vary with successional states and whether shift in plant functional group composition can explain the temporal variation in these responses. Previous studies at this experimental site showed a lack of climate change effect on productivity in a relative short term (Xu *et al.*, 2016). Taking advantage of all the eight-year data and a successional change in species composition, we show altered climate sensitivity of aboveground net primary productivity (ANPP) in the two distinct compositional states of plant functional groups. Specifically, we hypothesized that due to the temporal shift in functional group composition (i) the effects of precipitation and warming treatments on ANPP would change through time, (ii) the positive responses of ANPP to warming and double precipitation would amplify, and (iii) the negative response of ANPP to halved precipitation would dampen over time.

Materials and methods

Study site

The experiment was conducted in a grassland on the Kessler Atmospheric and Ecological Field Station in central Oklahoma, USA (34°59'N, 97°31'W). The study site is an abandoned

field from cropping. In 2008, we installed fencing around the experimental site to exclude large ungulate grazers; previously, this site had been grazed by cattle at moderate intensities for 40 years until 2008 (Xu *et al.*, 2016). The grassland was dominated by C₃ grass *Bromus japonicus* Thunb., C₃ forbs *Ambrosia trifida*, *Apocynum sp.*, *Solanum carolinense*, *Vernonia baldwinii*, *Securigera varia*, *Euphorbia dentate* and C₄ grass *Tridens flavus*, *Sorghum halepense*. Mean annual temperature was 16.3 °C with a monthly mean temperature of 4.4 °C in January and 27.7 °C in July, and mean annual precipitation is 914 mm (Oklahoma Climatological Survey, Norman, OK, USA). The soil texture class is loam with 51% of sand, 35% of silt and 13% of clay. The soil bulk density is 1.2 g cm⁻³. The concentrations of soil organic matter and total N are 1.9% and 0.1%, respectively.

Experimental design

The experiment used a paired factorial design with manipulation of air temperature and precipitation (Xu *et al.*, 2013, Xu *et al.*, 2016). There were two levels for warming: ambient temperature and elevated temperature, and three levels for precipitation: ambient precipitation, double precipitation (2 x ambient) and halved precipitation (1/2 ambient).

There were four random blocks, and within each there was one experimental plot (size 2.5 m × 3.5 m) for each factorial combination of warming and precipitation treatments ($N = 24$). For the warming treatment, the experimental plot was subjected to continuously warming by an infrared heater (100 W m⁻², Kalglo Electronics Inc, Bethlehem, PA, USA) hung 1.5 m above the ground in July 2009, while the other plot, equipped with a ‘dummy’ heater, served as a control. The distance between the warmed and control plots was approximately 5 m. For precipitation treatments, we applied a year-round rainfall-collection-redistribution device as described by Zhou *et al.* (2006) with the same area of the plot to double precipitation and a rainout-facility as described by Yahdjian and Sala (2002) to halve precipitation. In brief, the

rainout facility design is a fixed-location facility with a roof (3.66 m × 3 m) consisting of 18 strips (3.66m long, 8.25cm wide, 2.5mm thick) of transparent acrylic which can block about half of the rainfall while minimally affecting other environmental variables. The lower side of the rainout- facility tilts toward the prevailing upwind direction. The middle of the roof is 1.5m tall. Each 2.5 m × 3.5 m plot is divided into two 2.5 m × 1.75 m subplots. Plants were clipped in the southern 2.5 m × 1.75 m subplots which are at the opposite site of the prevailing upwind direction. Lateral water movement was prevented by inserting fiberglass sheet into the ground to a depth of 120 cm. In the control plots, we installed ‘dummy’ frames of the rain facility. Therefore, the six treatments are control (ambient) temperature and control precipitation (CC), control temperature and double precipitation (CD), control temperature and halved precipitation (CH), warming and control precipitation (WC), warming and double precipitation (WD), and warming and halved precipitation (WH).

Soil temperature, soil water content and precipitation measurements

Soil temperature was measured by thermocouples at a depth of 7.5 cm in the center of each plot. The thermocouples were connected to a CR10 datalogger (Campbell Scientific Inc., Logan, UT, USA) and recorded soil temperature every 15 min. Volumetric soil water content (v/v %) was measured once or twice a month using portable Time Domain Reflectometry equipment (Soil Moisture Equipment Corp., Santa Barbara, CA, USA) at a depth of 1-15cm. Precipitation data were obtained from an Oklahoma Mesonet Station (Washington Station) located approximately 200 m away from our experimental site.

Measurement of ANPP and its functional group biomass

ANPP, separated into C₃ and C₄ species, was directly measured by annually clipping at peak biomass (usually in September) in each plot. The clipped biomass was oven-dried at 70 °C for 48h and weighed to calculate ANPP and its function group biomass.

Statistical analyses

Before statistical analyses, the data were subjected to the Shapiro–Wilk test for normality and to the Levene’s test for homogeneity of variance. We used repeated-measures analysis of variance (ANOVA) to examine the main and interactive effects on soil temperature, soil water content, ANPP and its functional group biomass with warming and precipitation as main effects, year as the repeated factor, and block as a random effect. Multiple comparisons were conducted based on estimates of least square means to explore the difference among the three precipitation scenarios.

Linear regression was used to detect linear trends in C₃ and C₄ biomass, and to identify the relationship between C₃ and C₄ biomass. A general mixed-effect model was constructed with warming as the fixed factor and co-varying variables including annual mean air temperature, annual precipitation and annual mean functional composition (i.e., C₄ proportion) across time to explore the drivers of the inter-annual variability of the natural log response ratio (Ln rr) of ANPP to precipitation treatments. All statistical analyses were conducted in SAS v.8.1 (SAS Institute Inc., Cary, NC, USA).

Results

Environmental conditions

Mean annual air temperature over the experimental period (2009 ~ 2016) was 16.5°C with the range of 15.3 ~ 17.9 °C and mean annual precipitation was 911 mm with the range of 549 ~

1605mm (Fig. S1). There were no temporal trends in annual air temperature and annual precipitation over the eight-year experimental period (Fig. S1). Warming and precipitation treatments had substantial effects on soil temperature and water content (Fig. 1; Table S1). Eight-year warming increased soil temperature by 3°C on average and decreased soil water content by ca. 1.3% (absolute) with significant effect starting from 2013 (Fig. 1a; Table S1). Halved precipitation significantly increased soil temperature by 0.6 °C; precipitation treatments affected soil water content such that double precipitation increased soil water content by ca. 1% (absolute) and halved precipitation decreased soil water content by ca. 0.6% (absolute) with significant effect starting from 2012 (Fig. 1b; Table S1). Warming and precipitation treatments did not interact to affect soil temperature or soil water content.

Temporal change in functional group biomass and composition

From 2009 to 2016, C₃ biomass gradually decreased ($F_{1, 190} = 136.9$, $P < 0.0001$, $R^2 = 0.42$), and C₄ biomass increased ($F_{1, 190} = 37.45$, $P < 0.0001$, $R^2 = 0.16$) to compensate the loss of C₃ biomass in the control plots (Fig. 2a). These trends in functional group biomass were independent of warming (Fig. S2) or precipitation treatments (Fig. S3). An abrupt change in functional group composition occurred in 2014 (Fig. 2b). As a result, there were two distinct states of functional group composition through time: a C₃-dominated state from 2009 to 2013 (proportion of C₃ biomass: 71.0% on average over the five years) and a C₄-dominated state from 2014 to 2016 (proportion of C₄ biomass: 78.3% on average over the three years) (Fig. 2b). C₃ biomass is negatively associated with C₄ biomass in all the experimental plots ($F_{1, 190} = 77.67$, $P < 0.0001$, $R^2 = 0.29$; Fig. 3). A typical old-field successional change in species composition was associated with the functional group shift. The community transitioned from annual weedy grasses (e.g., *Bromus japonicus*) and annual forbs (e.g., *Ambrosia trifida*) to mostly perennial bunchgrass (e.g., *Tridens flavus* and *Sorghum halepense*).

Responses of total ANPP and functional group biomass to climate change

Given the compositional state shift in the two functional groups, we evaluated the effects of precipitation and warming treatments on ANPP and the functional group biomass (i.e., C₃ and C₄) in the two states within 2009 ~ 2013 and 2014 ~ 2016, respectively. Double precipitation did not affect ANPP in the first compositional state (C₃-dominated community), but increased ANPP by an average of 45.3% in the second compositional state (C₄-dominated community; Fig. 4a, b; Table 1). Halved precipitation reduced total ANPP in the first compositional state by an average of 17.6% yet did not affect ANPP in the second compositional state (Fig. 4c, d; Table 1). Warming did not affect ANPP in either of the two compositional states (Fig. 4e, f; Table 1). Furthermore, mixed-effect model showed that C₄% is a major factor accounting for the interannual variation in the natural log response ratio of ANPP (Ln rr) to altered precipitation (double precipitation: $F_{1,53} = 7.28$, $P = 0.009$; halved precipitation: $F_{1,53} = 3.02$, $P = 0.088$; Table 2).

We also examined how the two functional groups responded to climate change in the two compositional states. In the first state (2009 – 2013), double precipitation increased C₄ plant growth by 31.3% on average ($P = 0.036$) but did not affect C₃ biomass ($P = 0.19$; Fig. 5; Table 1); halved precipitation decreased C₃ biomass with a marginal significance by 21.7% on average ($P = 0.07$) but did not affect C₄ biomass ($P = 0.64$; Fig. 5; Table 1); warming did not influence either C₃ or C₄ biomass (Table 1). In the second state (2014-2016), double precipitation increased C₄ biomass by 69.6% on average ($P = 0.001$; Fig. 6a), but surprisingly reduced C₃ plant growth by 75.6% in the C₄-dominated community in the wettest year (i.e., 2015; Fig. 6b; Table 1); halved precipitation did not have an impact on either C₃ or C₄ biomass (Fig. 6b; Table 1); warming enhanced C₃ plant growth by 162% times in the wettest year (Fig. 6c; Table 1).

Discussion

Successional change in plant community

Our findings reveal the compositional state shift in the two plant functional groups over the eight experimental years. The studied temperate grassland transitioned from a C₃-dominant to a C₄-dominant system. The temporal trends in C₄ and C₃ biomass may be explained by the removal of the disturbance, i.e., grazing (Knapp & Medina, 1999, Koerner *et al.*, 2014). The recent enclosure in 2008 has kept the experimental site from herbivore grazing, which weakens the top down effects on the plant community (Koerner *et al.*, 2014, Post & Pedersen, 2008, Suttle *et al.*, 2007) and thus shifts the plant community from one state to another, a successional change. Specifically, C₃-dominated communities in the early state were mostly composed of annual forbs, including *Ambrosia trifida* (giant ragweed), *Solanum carolinense* (horsenettle) and *Euphorbia dentate* (toothed spurge), which were all weedy and generally unpalatable plant species. Removal of grazing released the dominant palatable C₄ grasses, including *Tridens flavus* (Purpletop) and the invasive *Sorghum halepense* (Johnson grass) from herbivore control with consequent spread.

The community shift from annual weedy grass (e.g., *Bromus japonicus*) and annual forbs (e.g., *Ambrosia trifida*) to mostly perennial bunch grass (e.g., *Tridens flavus* and *Sorghum halepense*) is consistent with other studies in grassland succession (Booth, 1941, Odum Eugene, 1960, Perino & Risser, 1972). For example, reduced foliage herbivory resulted in large increases in perennial grass growth and reduction in forb abundance (Brown & Gange, 1992). The fact that the same C₃ species are still dominating the plant community nearby under ambient conditions (i.e., grazed condition), and that an adjacent long-term experimental site which was fenced from grazing for over four decades features a C₄-dominated community (Shi *et al.*, 2016), indirectly supporting the grazing mechanism. The negative correlation between C₃ biomass and C₄ biomass further reveals possible antagonistic

interaction at the functional group level, which can explain the opposite temporal trends in C₃ versus C₄ biomass. Another mechanism could also account for the temporal trend in functional group is that the removal of grazing may reduce soil nitrogen availability (McNeil & Cushman, 2005) and lead to dominance of C₄ plant species, which can utilize nitrogen more efficiently than C₃ species (Lambers *et al.*, 1998). And it could also just be a release from grazers that preferentially choose the C₄ grasses over the less desirable C₃ forbs.

In addition, the abrupt shift from C₃ dominance to C₄ dominance during the study period happened in year 2014 which was extremely dry in terms of annual rainfall and soil water content. The sudden change in the compositional state suggests that the extreme dry year might have accelerated the rate of the successional change. This hypothesis is consistent with the result that halved precipitation treatment negatively affects the C₃ functional group. Besides dry year 2014, the wet years of 2015 and 2016 may have favored C₄ perennials over C₃ annuals. Without the extreme years, the plant community may have slowly converged to this C₄-dominated state, and this succession was sped up by an extreme event.

Responses of ANPP to climate change: long-term shift and associated mechanisms

We expected long-term shift in the responses of ANPP to climate change due to the documented transition from C₃- to C₄-dominated plant community (Langley & Megonigal, 2010, Morgan *et al.*, 2011, Zelikova *et al.*, 2014). Consistent with our prediction, amplified response of ANPP to double precipitation and dampened response of ANPP to halved precipitation were observed.

In the first compositional state (dominated by C₃ species), drought reduced ANPP through adversely influencing C₃ biomass and double precipitation increased C₄ biomass, as predicted by physiology. It could also be because C₃ species as annuals and C₄ species as perennials had differential sensitive to precipitation change. However, the increase in C₄ biomass was not proportionally enough to make a significant impact on the total ANPP. The

strong competition between C₃ and C₄ in double precipitation treatment may account for lack of response in ANPP to double precipitation in the first state. The findings are partially consistent with meta-analyses (Wilcox *et al.*, 2017, Wu *et al.*, 2011) which report positive and negative responses of ANPP to increased and decreased precipitation, respectively.

Altered sensitivity of ANPP to precipitation change in the second compositional state (dominated by C₄ species) in our study supports long-term shift in precipitation sensitivity (Smith *et al.*, 2009, Wilcox *et al.*, 2016). Double precipitation greatly increased the ANPP by enhancing C₄ biomass. Yet, halved precipitation did not reduce ANPP, which likely results from the well adaptations of the dominant C₄ plant species to dry conditions (Ehleringer *et al.*, 1997, Epstein *et al.*, 2002). In contrast to our expectation, halved precipitation did not affect C₃ biomass in the C₄-dominated community, possibly due to the fact that dry conditions may reduce interspecific competition (Kardol *et al.*, 2010) and alleviate the pressure on C₃ from C₄. Unexpected is also that increased precipitation reduced C₃ plant growth (Wilcox *et al.*, 2017, Wu *et al.*, 2011) in the C₄-dominated community in the wettest year (year 2015), likely due to the biotic competition with C₄ species that benefited from increased precipitation, highlighting the interactive nature of mechanisms that regulate climate sensitivity of ecosystem functions. In addition, C₃ species in this study are mostly annuals which are weak competitors compared to perennials.

We also predicted that the plant community in the second compositional state (C₄-dominated) would show greater positive response to warming given that C₄ plants are considered to be better adapted to warmer climates (Morgan *et al.*, 2011) than C₃ herbaceous plants. Instead, warming did not affect ANPP in either of the two states. The neutral response of ANPP to warming in the C₃-dominated community may be explained by the limited change in soil water content induced by warming and the relative insensitivity of C₃ plants in our studied community to warming; while the lack of response to warming and

warming-caused desiccation in the C₄-dominated community may be explained by the fact that C₄ species are well adapted to drought. This finding is consistent with the results from a semi-arid mixed-grass prairie showing that ANPP was unaffected by four years of warming (Morgan *et al.*, 2011). A similar finding was also reported in a old-field plant community, where warming did not affect the ANPP of the C₃-dominated system (Hoepfner & Dukes, 2012). In terms of individual responses of plant functional groups, warming did not affect C₃ or C₄ biomass in the first compositional state (C₃-dominate community). However, warming enhanced C₃ plant growth in the wettest year (2015) when the community was dominated by C₄ species. This supports that warming is likely to interact with extreme rainfall condition to exert impact on plant growth water availability (Jentsch *et al.*, 2007, Smith *et al.*, 2009).

Previous research has demonstrated various temporal trends in climate sensitivity of ecosystem functions. Amplified trends of soil C fluxes to warming was observed in both terrestrial (Xu *et al.*, 2015) and aquatic ecosystems (Yvon-Durocher *et al.*, 2017); attenuated trends of ecosystem functions such as (aboveground net primary productivity) ANPP and soil respiration to warming were also found in both grassland (Wu *et al.*, 2012) and forest ecosystems (Melillo *et al.*, 2002). So were the lack of temporal trends to climate change (Mueller *et al.*, 2016, Zelikova *et al.*, 2014, Zhu *et al.*, 2016). Results of this study showed all possible scenarios of the altered sensitivity of ecosystem productivity to long-term climate change: amplified sensitivity to increased precipitation, dampened sensitivity to decreased precipitation and lack of response to warming over time. In contrast to previous identified mechanisms, we found strong evidence that successional change in plant community was the contributing mechanism behind both the amplified and dampened responses.

Overall, the altered sensitivity of ANPP to precipitation change and the lack of response of ANPP to long-term warming highlight the predominant role of water availability in driving grassland ecosystem responses. The primary role of water – not temperature – is

consistent with a global climate sensitivity study, in which precipitation sensitivity is predominant in grassland ecosystems (Seddon *et al.*, 2016). In addition, the diverse responses of plant functional group biomass to climate change suggest that, besides plant physiology, there are other dominant factors such as biotic competition moderating long-term ecosystem responses, emphasizing the complexity of ecosystem responses to climate change.

Our findings have significant implications for understanding the linkage between plant community and ecosystem functioning in the context of long-term climate change. First, altered climate sensitivity with transition in the functional group composition highlights the importance of understanding the mechanisms underlying such compositional state shift; and the significance of involving vegetation dynamics in predicting future carbon state. Second, if climate change would affect species composition in the future (Cramer *et al.*, 2001, Ehleringer *et al.*, 1997, Epstein *et al.*, 2002), shift in species composition could in turn act as a long-term feedback to alter the ecosystem responses to climate change. In addition, long-term climate change experiments in early successional systems are essential for understanding the changes in strength and direction of ecosystem responses to climate change (Kroel-Dulay *et al.*, 2015).

Acknowledgements

We would like to thank many laboratory members for their help with field work. **Funding:** This study is financially supported by the United States Department of Energy, Biological Systems Research on the Role of Microbial Communities in Carbon Cycling Program (DE-SC0004601 and DE-SC0010715), by the US National Science Foundation (NSF) (DEB 0743778, DEB 0840964, DBI 0850290, and EPS 0919466), and by the Office of the Vice President for Research at the University of Oklahoma.

Reference

- Bardgett RD, Wardle DA (2003) HERBIVORE-MEDIATED LINKAGES BETWEEN ABOVEGROUND AND BELOWGROUND COMMUNITIES. *Ecology*, **84**, 2258-2268.
- Booth WE (1941) Revegetation of Abandoned Fields in Kansas and Oklahoma. *American Journal of Botany*, **28**, 415-422.
- Bradford MA, Davies CA, Frey SD *et al.* (2008) Thermal adaptation of soil microbial respiration to elevated temperature. *Ecology Letters*, **11**, 1316-1327.
- Brown VK, Gange AC (1992) Secondary plant succession: how is it modified by insect herbivory? *Vegetatio*, **101**, 3-13.
- Cramer W, Bondeau A, Woodward FI *et al.* (2001) Global response of terrestrial ecosystem structure and function to CO₂ and climate change: results from six dynamic global vegetation models. *Global Change Biology*, **7**, 357-373.
- Crowther TW, Bradford MA (2013) Thermal acclimation in widespread heterotrophic soil microbes. *Ecology Letters*, **16**, 469-477.
- Davidson DW (1993) The Effects of Herbivory and Granivory on Terrestrial Plant Succession. *Oikos*, **68**, 23-35.
- Ehleringer JR, Cerling TE, Helliker BR (1997) C₄ photosynthesis, atmospheric CO₂, and climate. *Oecologia*, **112**, 285-299.
- Ellery K, Ellery WN, Verhagen BT (1992) The distribution of C₃ and C₄ plants in a successional sequence in the Okavango Delta. *South African Journal of Botany*, **58**, 400-402.
- Epstein HE, Gill RA, Paruelo JM, Lauenroth WK, Jia GJ, Burke IC (2002) The relative abundance of three plant functional types in temperate grasslands and shrublands of North and South America: effects of projected climate change. *Journal of Biogeography*, **29**, 875-888.
- Fu YH, Zhao H, Piao S *et al.* (2015) Declining global warming effects on the phenology of spring leaf unfolding. *Nature*, **526**, 104-107.
- Grime JP, Brown VK, Thompson K *et al.* (2000) The response of two contrasting limestone grasslands to simulated climate change. *Science*, **289**, 762-765.
- Hoepfner SS, Dukes JS (2012) Interactive responses of old-field plant growth and composition to warming and precipitation. *Global Change Biology*, **18**, 1754-1768.

- Hoover DL, Knapp AK, Smith MD (2014) Resistance and resilience of a grassland ecosystem to climate extremes. *Ecology*, **95**, 2646-2656.
- Jentsch A, Kreyling J, Beierkuhnlein C (2007) A new generation of climate-change experiments: events, not trends. *Frontiers in Ecology and the Environment*, **5**, 365-374.
- Kardol P, Campany CE, Souza L, Norby RJ, Weltzin JF, Classen AT (2010) Climate change effects on plant biomass alter dominance patterns and community evenness in an experimental old-field ecosystem. *Global Change Biology*, **16**, 2676-2687.
- Knapp A, Medina E (1999) Success of C4 photosynthesis in the field: lessons from communities dominated by C4 plants. *C*, **4**, 251-283.
- Koerner SE, Burkepile DE, Fynn RWS *et al.* (2014) Plant community response to loss of large herbivores differs between North American and South African savanna grasslands. *Ecology*, **95**, 808-816.
- Kroel-Dulay G, Ransijn J, Schmidt IK *et al.* (2015) Increased sensitivity to climate change in disturbed ecosystems. *Nature Communications*, **6**, 10.1038/ncomms7682.
- Lambers H, Chapin FS, Pons TL (1998) Photosynthesis, Respiration, and Long-Distance Transport. In: *Plant Physiological Ecology*. pp Page. New York, NY, Springer New York.
- Langley JA, Megonigal JP (2010) Ecosystem response to elevated CO2 levels limited by nitrogen-induced plant species shift. *Nature*, **466**, 96-99.
- Luo YQ, Wan SQ, Hui DF, Wallace LL (2001) Acclimatization of soil respiration to warming in a tall grass prairie. *Nature*, **413**, 622-625.
- Mcneil SG, Cushman JH (2005) Indirect effects of deer herbivory on local nitrogen availability in a coastal dune ecosystem. *Oikos*, **110**, 124-132.
- Melillo JM, Steudler PA, Aber JD *et al.* (2002) Soil Warming and Carbon-Cycle Feedbacks to the Climate System. *Science*, **298**, 2173-2176.
- Morgan JA, Lecain DR, Pendall E *et al.* (2011) C-4 grasses prosper as carbon dioxide eliminates desiccation in warmed semi-arid grassland. *Nature*, **476**, 202-206.
- Mueller KE, Blumenthal DM, Pendall E *et al.* (2016) Impacts of warming and elevated CO2 on a semi-arid grassland are non-additive, shift with precipitation, and reverse over time. *Ecology Letters*, **19**, 956-966.
- Niu SL, Sherry RA, Zhou XH, Wan SQ, Luo YQ (2010) Nitrogen regulation of the climate-carbon feedback: evidence from a long-term global change experiment. *Ecology*, **91**, 3261-3273.

- Niu SL, Wan SQ (2008) Warming changes plant competitive hierarchy in a temperate steppe in northern China. *Journal of Plant Ecology*, **1**, 103-110.
- Odum EP (1969) The Strategy of Ecosystem Development. *Science*, **164**, 262-270.
- Odum Eugene P (1960) Organic Production and Turnover in Old Field Succession. *Ecology*, **41**, 34-49.
- Parmesan C (2007) Influences of species, latitudes and methodologies on estimates of phenological response to global warming. *Global Change Biology*, **13**, 1860-1872.
- Perino JV, Risser PG (1972) Some Aspects of Structure and Function in Oklahoma Old-Field Succession. *Bulletin of the Torrey Botanical Club*, **99**, 233-239.
- Piao S, Liu Z, Wang T *et al.* (2017) Weakening temperature control on the interannual variations of spring carbon uptake across northern lands. *Nature Clim. Change*, **7**, 359-363.
- Pickett STA, Collins SL, Armesto JJ (1987) Models, mechanisms and pathways of succession. *The Botanical Review*, **53**, 335-371.
- Post E, Pedersen C (2008) Opposing plant community responses to warming with and without herbivores. *Proceedings of the National Academy of Sciences*, **105**, 12353-12358.
- Reyes-Fox M, Steltzer H, Trlica MJ, McMaster GS, Andales AA, Lecain DR, Morgan JA (2014) Elevated CO₂ further lengthens growing season under warming conditions. *Nature*, **510**, 259-262.
- Seddon AWR, Macias-Fauria M, Long PR, Benz D, Willis KJ (2016) Sensitivity of global terrestrial ecosystems to climate variability. *Nature*, **531**, 229-232.
- Shi Z, Sherry R, Xu X *et al.* (2015) Evidence for long-term shift in plant community composition under decadal experimental warming. *Journal of Ecology*, **103**, 1131-1140.
- Shi Z, Xu X, Souza L *et al.* (2016) Dual mechanisms regulate ecosystem stability under decade-long warming and hay harvest. *Nature Communications*, **7**, 10.1038/ncomms11973.
- Smith MD, Knapp AK, Collins SL (2009) A framework for assessing ecosystem dynamics in response to chronic resource alterations induced by global change. *Ecology*, **90**, 3279-3289.
- Smith NG, Malyshev SL, Shevliakova E, Kattge J, Dukes JS (2016) Foliar temperature acclimation reduces simulated carbon sensitivity to climate. *Nature Clim. Change*, **6**, 407-411.

- Suttle KB, Thomsen MA, Power ME (2007) Species Interactions Reverse Grassland Responses to Changing Climate. *Science*, **315**, 640-642.
- Taylor SH, Ripley BS, Martin T, De-Wet L-A, Woodward FI, Osborne CP (2014) Physiological advantages of C4 grasses in the field: a comparative experiment demonstrating the importance of drought. *Global Change Biology*, **20**, 1992-2003.
- Von Fischer JC, Tieszen LL, Schimel DS (2008) Climate controls on C3 vs. C4 productivity in North American grasslands from carbon isotope composition of soil organic matter. *Global Change Biology*, **14**, 1141-1155.
- Wilcox KR, Blair JM, Smith MD, Knapp AK (2016) Does ecosystem sensitivity to precipitation at the site-level conform to regional-scale predictions? *Ecology*, **97**, 561-568.
- Wilcox KR, Shi Z, Gherardi LA *et al.* (2017) Asymmetric responses of primary productivity to precipitation extremes: A synthesis of grassland precipitation manipulation experiments. *Global Change Biology*, **23**, 4376-4385.
- Wilcox KR, Von Fischer JC, Muscha JM, Petersen MK, Knapp AK (2015) Contrasting above- and belowground sensitivity of three Great Plains grasslands to altered rainfall regimes. *Global Change Biology*, **21**, 335-344.
- Wu ZT, Dijkstra P, Koch GW, Hungate BA (2012) Biogeochemical and ecological feedbacks in grassland responses to warming. *Nature Climate Change*, **2**, 458-461.
- Wu ZT, Dijkstra P, Koch GW, Penuelas J, Hungate BA (2011) Responses of terrestrial ecosystems to temperature and precipitation change: a meta-analysis of experimental manipulation. *Global Change Biology*, **17**, 927-942.
- Xu X, Sherry RA, Niu S, Li D, Luo Y (2013) Net primary productivity and rain-use efficiency as affected by warming, altered precipitation, and clipping in a mixed-grass prairie. *Global Change Biology*, **19**, 2753-2764.
- Xu X, Shi Z, Chen X *et al.* (2016) Unchanged carbon balance driven by equivalent responses of production and respiration to climate change in a mixed-grass prairie. *Global Change Biology*, **22**, 1857-1866.
- Xu X, Shi Z, Li D, Zhou X, Sherry RA, Luo Y (2015) Plant community structure regulates responses of prairie soil respiration to decadal experimental warming. *Global Change Biology*, **21**, 3846-3853.
- Yahdjian L, Sala OE (2002) A rainout shelter design for intercepting different amounts of rainfall. *Oecologia*, **133**, 95-101.

Yvon-Durocher G, Hulatt CJ, Woodward G, Trimmer M (2017) Long-term warming amplifies shifts in the carbon cycle of experimental ponds. *Nature Clim. Change*, **7**, 209-213.

Zelikova TJ, Blumenthal DM, Williams DG, Souza L, Lecain DR, Morgan J, Pendall E (2014) Long-term exposure to elevated CO₂ enhances plant community stability by suppressing dominant plant species in a mixed-grass prairie. *Proceedings of the National Academy of Sciences of the United States of America*, **111**, 15456-15461.

Zhou X, Sherry RA, An Y, Wallace LL, Luo Y (2006) Main and interactive effects of warming, clipping, and doubled precipitation on soil CO₂ efflux in a grassland ecosystem. *Global Biogeochemical Cycles*, **20**, n/a-n/a.

Zhu K, Chiariello NR, Tobeck T, Fukami T, Field CB (2016) Nonlinear, interacting responses to climate limit grassland production under global change. *Proceedings of the National Academy of Sciences*, **113**, 10589-10594.

Table 1 Results (P values with F values in the brackets) of repeated-measures ANOVA for the responses of aboveground net primary productivity (ANPP), C₃ biomass and C₄ biomass to warming (W), altered precipitation (PPT), year and their interactions within 2009 ~ 2013 (C₃-dominated) and 2014 ~ 2016 (C₄-dominated), respectively. P values smaller than 0.05 are in bold.

	C ₃ -dominant				C ₄ -dominant			
	df	ANPP	C ₃	C ₄	df	ANPP	C ₃	C ₄
W	1, 15	0.903 (0.02)	0.382 (0.81)	0.386 (0.80)	1, 15	0.802 (0.06)	0.045 (4.77)	0.294 (1.18)
PPT	2, 15	0.011 (6.13)	0.175 (1.96)	0.031 (4.41)	2, 15	0.011 (6.23)	0.066 (3.28)	0.001 (11.18)
W × PPT	2, 15	0.193 (1.84)	0.133 (2.32)	0.622 (0.49)	2, 15	0.590 (0.55)	0.690 (0.38)	0.723 (0.33)
Year	4, 72	< 0.001 (11.35)	< 0.001 (18.32)	0.038 (2.69)	2, 36	< 0.001 (23.00)	< 0.001 (14.50)	< 0.001 (10.89)
W × Year	4, 72	0.516 (0.82)	0.726 (0.51)	0.643 (0.63)	2, 36	0.755 (0.28)	0.014 (4.86)	0.887 (0.12)
PPT × Year	8, 72	0.625 (0.78)	0.206 (1.41)	0.987 (0.22)	4, 36	0.913 (0.24)	0.024 (3.19)	0.684 (0.57)
W × PPT × Year	8, 72	0.850 (0.50)	0.401 (1.06)	0.876 (0.47)	4, 36	0.341 (1.17)	0.343 (1.16)	0.482 (0.89)

Table 2. Effects of warming (W), proportion of C₄ biomass (C₄%), annual precipitation (PPT), and annual mean temperature (T_{air}) on the natural log response ratio of ANPP (Ln rr) under double precipitation and halved precipitation treatments.

Effect	DF	F Value	Pr > F
<i>Ln rr, Double precipitation</i>			
W	1, 6	1.66	0.245
C ₄ %	1, 53	7.28	0.009
PPT	1, 53	1.03	0.315
T _{air}	1, 53	0.21	0.650
<i>Ln rr, Halved precipitation</i>			
W	1, 6	0.32	0.594
C ₄ %	1, 53	3.02	0.088
PPT	1, 53	0.03	0.872
T _{air}	1, 53	0.14	0.711

Figure captions

FIGURE 1. Responses of soil temperature (ST) and volumetric soil water content (SWC) to climate change within 2009 ~ 2016. Standard errors were omitted for clarity. Eight-year warming increased soil temperature by 3 °C on average, and halved precipitation significantly increased soil temperature by 0.6 °C (a). Warming decreased soil water content by ca. 1.3% (absolute) with significant effect starting from 2013. Precipitation treatments affected soil water content with double precipitation increasing soil water content by ca. 1% (absolute) and halved precipitation decreasing soil water content by ca. 0.6% (absolute) with significant effects starting from 2012 (b). Warming and precipitation change did not interact to affect soil temperature or soil water content. The six treatments are control (ambient) temperature and control precipitation (CC), control temperature and double precipitation (CD), control temperature and halved precipitation (CH), warming and control precipitation (WC), warming and double precipitation (WD), and warming and halved precipitation (WH). See Table S1 for statistics.

FIGURE 2 Temporal trends in functional group biomass and composition. C₃ biomass (open circles) decreased linearly over time ($F_{1,190} = 136.9$, $P < 0.0001$, $R^2 = 0.42$) and C₄ biomass (solid circles) increased linearly over time ($F_{1,190} = 37.45$, $P < 0.0001$, $R^2 = 0.16$); grey and black lines show linear fit with 95% confidence interval (a); C₃ (open circles) and C₄ proportion (solid circles) showed a drastic shift in 2014, with a sharp decrease in C₃ proportion and increase in C₄ proportion (b). Each point represents mean and standard error of the mean across all control experimental plots ($n = 4$).

FIGURE 3 The correlation between functional group biomass. The negative correlation between C₃ biomass and C₄ biomass ($F_{1, 190} = 77.67$, $P < 0.0001$, $R^2 = 0.29$) was observed.

The original data were square-rooted. Data in all treatments were included within 2009 ~ 2016 ($n = 192$). Grey line shows linear fit with 95% confidence interval.

FIGURE 4 Long-term shift in the natural log response ratio of ANPP to climate change.

Interannual variation in natural log response ratio (Ln rr) of ANPP to DP, double precipitation (**a**) showed significant difference between the two compositional states (**b**), interannual variation in Ln rr of ANPP to HP, halved precipitation (**c**) showed significant difference between the two compositional states (**d**), and interannual variation in Ln rr of ANPP to warming (**e**) showed no difference between the two compositional states (**f**). Each point represents mean and standard error of the mean across replicates ($n = 8$ for precipitation treatments and $n = 12$ for warming treatment). Note that the C₃-dominated community was within 2009 ~ 2013 and the C₄-dominated community was within 2014 ~ 2016. See Table 1 for statistical results.

FIGURE 5 Responses of total ANPP and biomass of the functional groups to climate change within 2009 ~ 2013. Different letters or numbers represent significant difference

among treatments at $\alpha = 0.05$. Uppercases are for C₃ biomass, lowercases are for C₄ biomass and numbers are for total ANPP. *P* values for the multiple comparisons in ANPP between the three precipitation treatments are (AMB vs DP: 0.25; AMB vs HP: 0.04; DP vs HP: 0.0036); comparisons in C₃ biomass (AMB vs DP: 0.19; AMB vs HP: 0.07; DP vs HP: 0.59); comparisons in C₄ biomass (AMB vs DP: 0.036; AMB vs HP: 0.64; DP vs HP: 0.014). AMB

is ambient precipitation, DP is double precipitation, and HP is halved precipitation. Gray bars are C₄ biomass, and black bars are C₃ biomass. See Table 1 for statistical results.

FIGURE 6 Responses of total ANPP and biomass of the functional groups to climate change within 2014 ~ 2016. *P* values for the multiple comparisons in ANPP between the three precipitation treatments are (AMB vs DP: 0.007; AMB vs HP: 0.89; DP vs HP: 0.0092); comparisons in C₄ biomass (AMB vs DP: 0.001; AMB vs HP: 0.96; DP vs HP: 0.0009). Double precipitation increased ANPP through positive effect on C₄ biomass (**a**, gray bars are C₄ biomass and black bars are C₃ biomass). In panel a, different letters or numbers represent significant difference among treatments at $\alpha = 0.05$. Lowercases are for C₄ biomass and numbers are for total ANPP. Double precipitation decreased C₃ biomass in 2015 (**b**) and warming increased C₃ biomass in 2015, the wettest year (**c**). In panel (**a**), AMB is ambient precipitation, DP is double precipitation, and HP is halved precipitation; in panel (**b**), solid circles represent halved precipitation, open circles are ambient precipitation, and triangles are double precipitation. In panel (**c**), open circles represent warming, and solid circle represent unwarmed treatment. See Table 1 for statistical results.











